

February 10, 2003

# UPGRADING THE THEOREM ON LOCAL ERGODICITY

Nándor Simányi<sup>1</sup>

University of Alabama at Birmingham  
 Department of Mathematics  
 Campbell Hall, Birmingham  
 AL 35294 U.S.A.  
 E-mail: simanyi@math.uab.edu

**Abstract.** We prove here that in the Theorem on Local Ergodicity for Semi-Dispersive Billiards (proved by N. I. Chernov and Ya. G. Sinai in 1987) the condition of the so called “Ansatz” can be dropped. That condition assumed that almost every singular phase point had a hyperbolic trajectory after the singularity. Having this condition dropped, the cited theorem becomes much stronger and easier to apply. At the end of the paper two immediate corollaries of this improvement are discussed: One of them is the (fully hyperbolic) Bernoulli mixing property of every hard disk system ( $D = 2$ ), the other one claims that the ergodic components of every hard ball system ( $D \geq 3$ ) are open.

Primary subject classification: 37D50

Secondary subject classification: 34D05

---

<sup>1</sup>Research supported by the National Science Foundation, grant DMS-0098773.

## §1. INTRODUCTION

**Semi-Dispersive Billiards.** Assume that  $\mathbf{Q} \subset \mathbb{R}^d$  or  $\mathbf{Q} \subset \mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$  ( $d \geq 2$ ) is an open and connected subset with a non-empty, piecewise smooth (i. e. piecewise  $C^\infty$ ) boundary  $\partial\mathbf{Q}$  and compact closure  $\overline{\mathbf{Q}} = \mathbf{Q} \cup \partial\mathbf{Q}$ . In order to avoid unnecessary technical complications, we also assume that the  $d$ -dimensional spatial angle  $\angle(q)$  subtended by  $\mathbf{Q}$  at any of its boundary points  $q \in \partial\mathbf{Q}$  is positive. We also assume that the smooth components  $\partial\mathbf{Q}_i$  ( $i = 1, 2, \dots, k$ ) of  $\partial\mathbf{Q}$  are (not necessarily strictly) concave, i. e. they are “bending away” from  $\mathbf{Q}$ . In technical terms this means the following: If we supply the smooth component  $\partial\mathbf{Q}_i$  with the field  $n(q)$  ( $q \in \partial\mathbf{Q}_i$ ) of unit normal vectors pointing inward  $\mathbf{Q}$ , then the second fundamental form  $K(q)$  of  $\partial\mathbf{Q}_i$  at  $q$  (with respect to the selected field of unit normal vectors) is a non-negative operator for any  $q \in \partial\mathbf{Q}_i$  and  $i = 1, \dots, k$ . Finally, one always assumes that at any boundary point  $q \in \partial\mathbf{Q}_i \cap \partial\mathbf{Q}_j$  ( $1 \leq i < j \leq k$ ) the tangent vectors of  $\mathbb{R}^d$  (or  $\mathbb{T}^d$ ) at  $q$  pointing inward  $\mathbf{Q}$  form a convex cone.

The semi-dispersive billiard system (flow)  $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$  is a dynamical system describing the uniform motion (with unit speed) of a point particle inside the flat domain  $\mathbf{Q}$  that bounces back at  $\partial\mathbf{Q}$  in a fully elastic manner, according to the law of geometric optics, that is, the angle of reflection equals the angle of incidence. The phase space  $\mathbf{M}$  of the arising flow is essentially the unit tangent bundle  $\overline{\mathbf{Q}} \times \mathbb{S}^{d-1}$  ( $\mathbb{S}^{d-1}$  being the unit sphere of velocities) modulo the natural identification of the incoming (pre-collision) and outgoing (post-collision) velocities at the boundary  $\partial\mathbf{M} = \partial\mathbf{Q} \times \mathbb{S}^{d-1}$ . The flow  $\{S^t\}_{t \in \mathbb{R}}$  is the time-evolution of the system, and the (invariant) Liouville measure  $\mu$  turns out to be the normalized Lebesgue measure:  $d\mu = \text{const} \cdot dq \cdot dv$ , where  $x = (q, v) \in \mathbf{M}$ ,  $q \in \mathbf{Q}$ ,  $v \in \mathbb{S}^{d-1}$  is the standard expansion of a phase point  $x$  into the configuration and velocity components.

In the case  $K(q) > 0$  (for every  $q \in \partial\mathbf{Q}$ ) we speak about a “dispersive billiard system”.

The primary examples of semi-dispersive billiards are the so called cylindric billiards (see [S-Sz(2000)]), among which the most important special models are the hard ball systems, see §2 of [S-Sz(1999)]. For a more detailed introduction to these dynamical systems, please see §2 (the sections containing the prerequisites) of the papers [K-S-Sz(1990-A)], [S-Sz(1999)], and [Sim(2002)].

Assume that  $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$  is a semi-dispersive billiard flow with the following additional properties:

(1) the smooth components of  $\partial\mathbf{Q}$  are algebraic (i. e. polynomially defined) hyper-surfaces;

(2) for  $\nu$ -almost every singular phase point  $x \in \mathcal{SR}^+ (\subset \partial\mathbf{M})$  the forward semi-trajectory  $S^{[0, \infty)}x$  of  $x$  is sufficient (or *geometrically hyperbolic*, see §2);

(3) a given phase point  $x_0 \in \mathbf{M} \setminus \partial\mathbf{M}$  has a sufficient orbit  $S^{\mathbb{R}}x_0$  (see §2) with at most one (and simple) singularity on it.

Under the above conditions, the Theorem on Local Ergodicity claims that some open neighborhood  $U_0 \ni x_0$  of  $x_0$  belongs (modulo the zero sets) to a single ergodic component of the flow  $\{S^t\}_{t \in \mathbb{R}}$ .

This fundamental result was originally proved by N. I. Chernov and Ya. G. Sinai as Theorem 5 in [S-Ch(1987)], being actually the corollary of the rather technical Lemma 3 there. Later on, the proof of Lemma 3 was somewhat clarified and put into a more general context as Theorem 3.6 of [K-S-Sz(1990-A)]. In that paper the exact counterpart of the above cited Theorem on Local Ergodicity is Corollary 3.12. In these two papers the above condition (1) (i. e. the “algebraicity”) was not yet present.

It turned out recently that the proof, as a matter of fact, requires this condition, see [B-Ch-Sz-T(2002)], where the main result is listed as Theorem 4.4 (the analogue of the technical Lemma 3 mentioned above). It was just the paper [B-Ch-Sz-T(2002)] in which the authors noticed the necessity to include the algebraicity condition (1) and, actually, they fixed the flaw of the earlier proofs.

Condition (2) is often called the “Chernov-Sinai Ansatz” in the literature.

Our goal here is to show the reader that the Ansatz (Condition (2)) may be dropped from the Theorem on Local Ergodicity altogether without hurting its validity! We will be doing so by introducing some interesting (but not too big) changes in the original proof of Theorem 3.6 of [K-S-Sz(1990-A)]. Our main reference will be just that paper.

## §2. PREREQUISITES

**Singularities and Trajectory Branches.** The billiard system  $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$  has two types of singularities. The first one is caused by the presence of the so called “tangential reflections”  $x_0 = (q_0, v_0) \in \partial\mathbf{M}$  for which  $\langle v_0, n(q_0) \rangle = 0$ , i. e. the velocity  $v_0$  happens to lie inside the tangent hyperplane  $\mathcal{T}_{q_0}\partial\mathbf{Q}$  of  $\partial\mathbf{Q}$  at  $q_0 \in \partial\mathbf{Q}$ . If the trajectory  $S^{(-\infty, \infty)}y$  of a phase point hits a tangential reflection at time  $t_0 > 0$  ( $S^{t_0}y = x_0$ ), then the billiard map  $S^{t_0+\epsilon}$  is still continuous at  $y$ , though it ceases to be smooth. The often studied first return map  $T : \partial\mathbf{M} \rightarrow \partial\mathbf{M}$  of the boundary  $\partial\mathbf{M}$  is no longer even continuous at the inverse image  $T^{-1}x_0$  of a phase point  $x_0$  with tangential reflection.

The second type of singularity, the so called “multiple reflection” takes place when the flow  $S^{t_0}y = x_0 = (q_0, v_0^-) \in \partial\mathbf{M}$  ( $v_0^-$  is the “incoming” velocity) hits more than one boundary components of  $\mathbf{M}$ , i. e. when  $q_0 \in \partial\mathbf{Q}_i \cap \partial\mathbf{Q}_j$  with

$1 \leq i < j \leq k$ . We are going to briefly describe the discontinuity of the flow  $\{S^t\}$  caused by a multiple collision at time  $t_0$ . Assume first that the pre-collision velocity is given. What can we say about the possible post-collision velocity? Let us perturb the pre-collision phase point (at time  $t_0 - 0$ ) infinitesimally, so that the collisions at  $\sim t_0$  occur at infinitesimally different moments. By applying the collision laws to the arising finite sequence of collisions, (the finiteness follows from Theorem 1 of [B-F-K(1998)]) we see that on the two sides of the current (codimension-one) singularity manifold there are two, significantly different continuations (the so called branches) of the trajectory: On one side the finite sequence of collisions near time  $t_0$  begins with a  $\partial\mathbf{Q}_i$ -reflection, then it continues in an alternating manner with  $\partial\mathbf{Q}_j$ ,  $\partial\mathbf{Q}_i, \dots$  reflections all the way until the resulting (reflected) velocity  $v^+$  happens to point inward  $\mathbf{Q}$ . On the other side of the singularity, however, the corresponding finite, alternating sequence of reflections begins with a  $\partial\mathbf{Q}_j$ -collision. These two continuations (branches) result in two (typically different) outgoing velocities  $v^+$ . Analogous statements can be made about the backward continuations and the  $v^+ \mapsto v^-$  correspondence. It follows from Lemma 4.1 of [K-S-Sz(1990-A)] that a typical singular phase point  $x_0$  has the property that its trajectory hits a singularity only once, say, at time  $t_0 > 0$  and

- (a)  $q(S^{t_0}x_0)$  lies only on one boundary component  $\partial\mathbf{Q}_i$  if  $S^{t_0}x_0$  is a tangential reflection;
- (b)  $q(S^{t_0}x_0)$  lies exactly on two boundary components  $\partial\mathbf{Q}_i$  and  $\partial\mathbf{Q}_j$  ( $i \neq j$ ) if  $S^{t_0}x_0$  is not a tangency;
- (c) none of the arising two trajectory branches hits a singularity any more.

In this case we say that the trajectory of  $x_0$  encounters a so called *simple singularity*. In this situation the entire orbit  $S^{(-\infty, \infty)}x_0$  has exactly two branches.

Since, in the case of multiple collisions, there is no unique continuation of the trajectories, we need to make a clear distinction between the set of reflections  $\mathcal{SR}^+$  supplied with the outgoing velocity  $v^+$ , and the set of reflections  $\mathcal{SR}^-$  supplied with the incoming velocity  $v^-$ . For typical phase points  $x^+ \in \mathcal{SR}^+$  the forward trajectory  $S^{[0, \infty)}x^+$  is non-singular and uniquely defined, and analogous statement holds true for typical phase points  $x^- \in \mathcal{SR}^-$  and the backward trajectory  $S^{(-\infty, 0]}x^-$ . For a more detailed exposition of singularities and trajectory branches, the reader is kindly referred to §2 of [K-S-Sz(1990-A)]. We denote by  $\mathcal{SR} = \mathcal{SR}^- \cup \mathcal{SR}^+$  the set of all singular reflections.

Finally, we note that the trajectory of the phase point  $x_0$  has exactly two branches, provided that  $S^t x_0$  hits a singularity for a single value  $t = t_0$ , and the phase point  $S^{t_0}x_0$  does not lie on the intersection of more than one singularity manifolds. (In this case we say that the trajectory of  $x_0$  has a “simple singularity”.)

**Sufficiency (Geometric Hyperbolicity).** Let  $S^{[a,b]}x_0$  be a non-singular (i. e. smooth) trajectory segment of a semi-dispersive billiard flow  $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$ . At the starting phase point  $y_0 = S^a x_0 = (q_0, v_0)$  we construct the submanifold

$$\Sigma_0 = \{(q, v_0) \mid q - q_0 \perp v_0 \text{ and } \|q - q_0\| < \epsilon_0\}$$

of  $\mathbf{M}$  with a fixed number  $\epsilon_0 \ll 1$ . Denote by  $\Sigma_1$  the connected component of the image  $S^{b-a}(\Sigma_0)$  containing  $S^b x_0 = S^{b-a} y_0$ . It is easy to see that the smooth submanifold  $\Sigma_1$  of  $\mathbf{M}$  has the form

$$\Sigma_1 = \{(q, v(q)) \mid q \in \tilde{\Sigma}_1\},$$

where  $\tilde{\Sigma}_1 \subset \mathbf{Q}$  is a smooth hypersurface of the configuration space  $\mathbf{Q}$  supplied with the field  $v(q)$  of unit normal vectors. A direct consequence of the semi-dispersive property is that the second fundamental form  $K(q)$  of  $\tilde{\Sigma}_1$  at any  $q \in \tilde{\Sigma}_1$  (with respect to the above field of unit normal vectors) is non-negative. The trajectory segment  $S^{[a,b]}x_0$  is said to be *sufficient* (or *geometrically hyperbolic*) if  $K(q_1) > 0$ , where  $S^b x_0 = (q_1, v_1)$ .

**Definition.** The trajectory segment  $S^{[a,b]}x$  containing exactly one singularity (a so called “simple singularity”, see above) is said to be sufficient if and only if both branches of this trajectory segment are sufficient.

**Definition.** The phase point  $x \in \mathbf{M}$  with at most one singularity is said to be sufficient if and only if its whole trajectory  $S^{(-\infty, \infty)}x$  is sufficient, which means, by definition, that some of its bounded segments  $S^{[a,b]}x$  are sufficient.

In the case of an orbit  $S^{(-\infty, \infty)}x$  with a single singularity, sufficiency means that both branches of  $S^{(-\infty, \infty)}x$  are sufficient.

The great importance of the sufficiency of a phase point  $x_0$  is given by the fact that (for the discrete time billiard map, i. e. for the first return map  $T$  to  $\partial\mathbf{M}$ ) such a phase point always has a suitably small, open neighborhood  $U_0 \ni x_0$  for which the first return map of  $T$  to  $U_0$  exhibits a uniformly hyperbolic behavior, see Lemma 2.13 in [K-S-Sz(1990-A)].

For a more detailed exposition of the above notions and facts, the reader is kindly referred to §2 of [K-S-Sz(1990-A)].

**No accumulation (of collisions) in finite time.** By the results of Vaserstein [V(1979)], Galperin [G(1981)] and Burago-Ferleger-Kononenko [B-F-K(1998)], in a semi-dispersive billiard flow there can only be finitely many collisions in finite time intervals, see Theorem 1 in [B-F-K(1998)]. Thus, the dynamics is well defined as long as the trajectory does not hit more than one boundary components at the same time.

### §3. THE IMPROVEMENTS

Let us see — step by step — all changes in the proof of Theorem 3.6 (of [K-S-Sz(1990-A)]) that facilitate the dropping of the Ansatz from the set of conditions for the Theorem on Local Ergodicity.

1. First of all, throughout the entire proof of Theorem 3.6 the distance function  $z(x)$  (measuring the distance between a phase point  $x \in \mathbf{M}$  and the set  $\mathcal{SR}$  of singular reflections) has to be replaced by the “tubular distance function”  $z_{\text{tub}}(x)$ , along the lines of 4. in the Erratum to the paper [K-S-Sz(1990-A)], i. e. in [K-S-Sz(1990-B)].

2. Concerning Definition 5.1 in the proof: The definition of the expansion coefficient

$$\kappa_{n,0}(y) = \left\| (D_{-T^n y, \Sigma}^n)^{-1} \right\|^{-1}$$

should not contain any infimum taken with respect to  $\Sigma$ . Instead, the orthogonal manifold  $\Sigma$  has to be the *unique* family of rays emanating from the fixed configuration point  $q(-T^n y) = q(T^n y)$  and containing *all* small velocity perturbations of the phase point  $-T^n y$ . (A so called “candle”, emanating light from one point in every (nearby) direction.) We note here that the natural metric on the “candle”  $\Sigma$  (needed in the use of the norm  $\| \cdot \|$  in the definition of  $\kappa_{n,0}(y)$ ) is the angle between the different velocities.

3. Regarding the definition of the expansion coefficient  $\kappa_{n,\delta}(y)$  in 5.1:

$$\kappa_{n,\delta}(y) = \inf_{w \in \Sigma} \left\| (D_{w, \Sigma}^n)^{-1} \right\|^{-1},$$

where  $\Sigma$  is again any “candle” manifold containing  $-T^n y$  (just as in the definition of  $\kappa_{n,0}(y)$ ) with the additional properties

- (i)  $T^n$  is smooth on  $\Sigma$ ,
- (ii)  $T^n \Sigma \subset B_\delta(-y)$ .

4. Lemma 5.3 is not used; should be dropped altogether. It is actually false for the re-defined expansion coefficient  $\kappa_{n,\delta}(y)$ . The only monotonicity property of these coefficients that is used in the proof pops up in the sentence beginning with “Finally, since  $\kappa_{n,\delta}(T^{-n}w)$  increases in  $n$ ” in the lower third of page 557 of the proof, and that monotonicity property is still obviously true.

5. Concerning the crucial construction of invariant manifolds, i. e. the proof of Lemma 5.4: The manifold  $\Sigma_n^n(y)$  should be the “candle” manifold (defined in 2. above) containing all rays emanating from the point  $q(-T^n y)$  and being closer

than  $c_3\delta$  to the phase point  $-T^n y$  (measured in the angular metric between the velocities). Then

$$\Sigma_0^n(y) = T^n \Sigma_n^n(y),$$

and  $-\tilde{\Sigma}_0^n = -\tilde{\Sigma}_0^n(y)$  is the connected component of the intersection

$$(-\Sigma_0^n(y)) \cap B_{c_3\delta}(y)$$

containing  $y$ , just as in the original proof. Otherwise, the proof of Lemma 5.4 remains intact.

6. It is worth reviewing the closing argument of the proof of Lemma 6.3, i. e. the four-line paragraph right after (6.6). Namely, if

$$\Sigma = \Sigma_{n_2}^{n_2}(T^{-n_2}w)$$

is a permitted “candle” manifold in the definition of  $\kappa_{n_2, c_3\delta}(T^{-n_2}w)$ , then  $-T^{n_1}\Sigma \subset U$ , since it contains the phase point  $T^{-n_1}w$ , and we may assume that this latter phase point is in a shrunk base neighborhood  $U_0 \subset U$  so that the entire manifold  $-T^{n_1}\Sigma$  with radius  $< c_3\delta$  is contained by  $U$ . Then the application of  $T^{n_2-n_1}$  to the manifold  $T^{n_1}\Sigma$  further expands this latter manifold by at least a factor of  $\Lambda$ , for  $-T^{n_2}\Sigma \subset U$ . However, the manifold  $\Sigma$  is obviously permitted in the definition of  $\kappa_{n_1, c_3\delta}(T^{-n_1}w)$ , so the expansion rate between  $\Sigma$  and  $T^{n_1}\Sigma$  is already at least  $\Lambda^m$ , thus the total expansion rate between  $\Sigma$  and  $T^{n_2}\Sigma$  becomes at least  $\Lambda^{m+1}$ , contradicting to (6.6).

7. The crucial change in the revised proof can be seen at the limit relation (6.8). The point is that, due to the new definition of the expansion coefficient  $\kappa_{n,0}(T^{-n}y)$ , the limit relation  $\lim_{n \rightarrow \infty} \kappa_{n,0}(T^{-n}y) = \infty$  of (6.8) now holds for *every* phase point  $y \in \mathcal{R}$ , irrespective of the Ansatz! Indeed, the coefficient  $\kappa_{n,0}(T^{-n}y)$  increases at least linearly in the absolute value  $|t|$  of the time  $t$  measured along the past trajectory  $\{S^t y \mid t < 0\}$ , as one easily sees by using the semi-dispersive characteristic of the billiard flow. However, according to Theorem 1 of [B-F-K(1998)], the time moments of collisions on any trajectory can not accumulate at any finite point. Therefore  $t \rightarrow -\infty$ , as  $n \rightarrow \infty$  in the above situation.

8. In the set inequality (6.10) one has to write

$$\bigcup_{n \geq N_\eta} T^n U_{n,m}^b \subset \mathcal{R}^{[c_3\delta]} \setminus \bigcup_{i=1}^l V_2(y_i, m)$$

in order to obtain a “regularly shaped” set  $\bigcup_{i=1}^l V_2(y_i, m)$  (instead of  $K_\eta$ ) and gain the important upper estimation  $\text{const} \cdot \eta$  for the measure of the set

$$\mathcal{R}^{[c_3\delta]} \setminus \bigcup_{i=1}^l V_2(y_i, m).$$

9. Finally, the “algebraicity” condition (2) in the Theorem on Local Ergodicity has to be assumed in order to save the proofs of the important geometric inequalities of Lemma 4.6, (5.11), and (5.12). The much needed fix for the original proof (in the algebraic case) came recently as the main result of the paper [B-Ch-Sz-T(2002)]. Fortunately, every cylindric billiard system, esp. every hard sphere system, is algebraic by nature.

## COROLLARIES

**Corollary 1.** The revised Theorem on Local Ergodicity, along with the Theorem of [Sim(2002)] gives us that *every* hard sphere system (in any dimension  $\nu \geq 2$ ) has *open* ergodic components and it is fully hyperbolic (i. e. has non-zero relevant Lyapunov exponents almost everywhere).

**Corollary 2.** The main result of the paper [Sim(2001)] is that almost every (in terms of the masses and the radius of the disks) hard disk system on the unit 2-torus  $\mathbb{T}^2$  is ergodic and fully hyperbolic. (These systems are actually Bernoulli flows.) As remark 9.3 of the cited paper explains, the dropping of a zero-measured family of hard disk systems (the annoying presence of the phrase “almost every” in the result) is necessitated merely by the cumbersome proof of the Ansatz. Having the above improvement of the Theorem on Local Ergodicity at hand, now we have a significantly stronger result: Every hard disk system  $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$  on  $\mathbb{T}^2$  with masses  $m_1, \dots, m_N$  and radius  $r$  is a Bernoulli flow and fully hyperbolic, of course, after having made the trivial reductions of the obviously preserved physical quantities.

## REFERENCES

- [B-Ch-Sz-T(2002)] P. Bálint, N. Chernov, D. Szász, I. P. Tóth, *Multidimensional semidispersing billiards: singularities and the fundamental theorem*, Ann. Henri Poincaré **3** no. 3, 451–482.
- [B-F-K(1998)] D. Burago, S. Ferleger, A. Kononenko, *A geometric approach to semi-dispersing billiards*, Ergod. Th. & Dynam. Sys. **18**, 303–319.
- [G(1981)] G. Galperin, *On systems of locally interacting and repelling particles moving in space*, Trudy MMO **43**, 142–196.
- [K-S-Sz(1990-A)] A. Krámli, N. Simányi, D. Szász, *A “Transversal” Fundamental Theorem for Semi-Dispersing Billiards*, Commun. Math. Phys. **129**, 535–560.



- [K-S-Sz(1990-B)] A. Krámli, N. Simányi, D. Szász, *Erratum. A “Transversal” Fundamental Theorem for Semi-Dispersing Billiards*, Commun. Math. Phys. **138**, 207–208.
- [Sim(2001)] N. Simányi, *Proof of the Boltzmann-Sinai Ergodic Hypothesis for Typical Hard Disk Systems*, Manuscript, submitted for publication. arXiv:math.DS/0008241.
- [Sim(2002)] N. Simányi, *The Complete Hyperbolicity of Cylindric Billiards*, Ergod. Th. & Dynam. Sys. **22**, 281-302, arXiv:math.DS/9906139.
- [S-Ch(1987)] Ya. G. Sinai, N.I. Chernov, *Ergodic properties of certain systems of 2-D discs and 3-D balls*, Russian Math. Surveys **(3) 42**, 181-207.
- [S-Sz(1999)] N. Simányi, D. Szász, *Hard Ball Systems Are Completely Hyperbolic*, Annals of Math. **149**, 35-96, arXiv:math.DS/9704229..
- [S-Sz(2000)] N. Simányi, D. Szász, *Non-integrability of Cylindric Billiards and Transitive Lie Group Actions*, Ergod. Th. & Dynam. Sys. **20**, 593-610.
- [V(1979)] L. N. Vaserstein, *On Systems of Particles with Finite Range and/or Repulsive Interactions*, Commun. Math. Phys. **69**, 31-56.